



Health literacy: a new piece of the puzzle in psoriasis care? A cross-sectional study*

M.H. Larsen ,¹ Y.A.S. Strumse,² C.R. Borge,^{1,3} R. Osborne ,⁴ M.H. Andersen^{1,5} and A.K. Wahl ¹

¹Department of Health Sciences, Institute of Health and Society, Faculty of Medicine, University of Oslo, Oslo, Norway

²Section for Climate Therapy and ⁵Department of Transplantation Medicine, Oslo University Hospital, Oslo, Norway

³Department of Patient Safety and Research at Lovisenberg Diaconal Hospital, Oslo, Norway

⁴Deakin University, Faculty of Health, School of Health and Social Development, Geelong, VIC, Australia

Linked Comment: Rowlands. *Br J Dermatol* 2019; **180**:1299–1300.

Summary

Correspondence

Marie H. Larsen.

E-mail: m.h.larsen@medisin.uio.no.

Accepted for publication

28 December 2018

Funding sources

University of Oslo, Norway. R.O. was funded, in part, through a National Health and Medical Research Council of Australia Senior Research Fellowship (#APP1059122).

*Plain language summary available online

DOI 10.1111/bjd.17595

Background Health literacy (HL) – the ability to seek, understand and utilize health information – is important for good health. Suboptimal HL has been associated with poorer health outcomes in other chronic conditions, although this has not previously been studied in patients with psoriasis.

Objectives To investigate the HL strengths and weaknesses of a cohort of patients with moderate-to-severe psoriasis. Another aim was to examine possible associations between patients' quality of life, their demographic, clinical and self-management characteristics, and dimensions of HL.

Methods A cross-sectional study was conducted. Data were collected from a cohort of patients with psoriasis who had received climate helio therapy from 2011 to 2016 (n = 825). HL was assessed by the Health Literacy Questionnaire (HLQ). The association between HL domains, demographic, clinical and self-management variables were analysed using bivariate correlation and a four-step linear multiple regression model.

Results The scores on all HLQ dimensions indicated lower health literacy than other populations. The linear regression models showed a significant association between HL, quality of life and self-management variables, with higher HL predicting higher quality of life, self-efficacy and psoriasis knowledge. Sex, educational attainment, age and disease severity had less influence on health literacy.

Conclusions Improving HL may be a useful strategy for reducing disparities in self-management skills for patients with psoriasis. Interventions that aim to reduce disease severity and increase psoriasis knowledge, self-efficacy and quality of life may positively increase HL.

What's already known about this topic?

- Health literacy (HL) has increasingly been acknowledged as an essential component of an individual's resources and competency to self-manage chronic conditions such as psoriasis.
- HL negatively affects healthcare utilization.
- HL has not yet been investigated in patients with psoriasis.

What does this study add?

- This is the first comprehensive HL assessment of psoriasis.
- Patients with psoriasis who have received climate helio therapy have lower HL scores than people with other chronic conditions.

- Factors associated with lower HL in people with psoriasis include lower educational attainment, lower psoriasis severity, lack of knowledge, lower self-efficacy, fewer comorbidities, shorter disease duration and lower quality of life.
- Assessing HL in psoriasis may help tailor self-management support interventions.

The worldwide prevalence of psoriasis ranges between 0.09% and 11.9%,^{1,2} making psoriasis a substantial global problem. Psoriasis leads to disfigurement, disability and marked loss of productivity, according to the World Health Organization's global report on psoriasis.³ There is also a significant negative impact on mental well-being, including higher rates of depression and other burdens across the lifespan, and on society overall.^{4–7} Thus, for many, a life with psoriasis includes physical, psychological and economic challenges.^{6,8–11} The cumulative impact of comorbidity risks related to psoriasis,^{12,13} in addition to time-consuming skin treatments, may also make treatment concordance and self-management difficult for patients.¹⁴ Given this biopsychosocial complexity and that of modern healthcare systems, the demands on these individuals in terms of self-management, healthcare system navigation and communication with different healthcare professionals appear to have increased.^{5,15,16}

Of the numerous biopsychosocial factors that may explain poor self-management in psoriasis, health literacy (HL) – defined as an individual's capacity to seek, understand and utilize health-information¹⁷ – has not yet been investigated. Suboptimal HL in people with chronic conditions such as asthma, diabetes and chronic obstructive pulmonary disease is reported to be associated with limited disease knowledge, poor self-management skills, limited understanding of health information and lower engagement with healthcare providers.^{18–23} These factors also appear to be important in the management of psoriasis, signifying that an investigation of HL among individuals with psoriasis is warranted.

Seeking treatment for psoriasis may be complex. Individuals may consult several healthcare professionals, consider multiple treatment options and receive large amounts of information on psoriasis, treatment, comorbidities and lifestyle recommendations.^{24,25} The latter might be contradictory and delivered in a manner that the individual cannot easily understand. Patients' general knowledge of psoriasis is known to be low,²⁶ especially knowledge related to the role of comorbidities.^{27,28} Efficacious psoriasis care may require HL skills such as collaborative communication with healthcare providers, taking an active part in shared-decision making, and participation in self-management and goal setting.^{16,29–33} However, people with HL weaknesses may lack the skills to accomplish such tasks and find it difficult to access and understand healthcare information and implement the recommended behaviours.

Conceptually, adequate health literacy in the context of psoriasis includes a constellation of skills that are critical for patients to manage their condition and navigate the healthcare environment effectively. To the best of our knowledge, there

have been no specific HL studies of people with psoriasis, certainly none that has used the recently developed comprehensive HL measures, which include HL related to critical appraisal, social and communication skills.³⁴ Hence, the aim of the present study was to describe the HL of people with psoriasis using the Health Literacy Questionnaire (HLQ), a robust nine-dimension assessment tool. The study sought to investigate associations between HL and demographic-, clinical- and self-management-related variables (e.g. self-efficacy and psoriasis knowledge) and quality of life in a cohort of patients with moderate-to-severe psoriasis.

Patients and methods

This paper is based on data from a cross-sectional study with a cohort of 1275 adults who all had previously (once or several times from 2011 to 2017) participated in the Norwegian climate helio therapy (CHT) programme in Gran Canaria. Norwegian patients with psoriasis have been offered CHT since 1976 and the cost of the therapy is covered by the Norwegian health authorities. Climate therapy is offered to patients who may require hospitalization and frequent intensive outpatient care. CHT may also be part of a sequential or tailored treatment regimen for many patients.³⁵ Hence, CHT is a routine part of Norwegian psoriasis care. All participants aged 18–85 years who had previously participated in CHT were invited by post to participate and were sent the information, the consent form and the survey questionnaire, including questions on health and demographics, to their home address. A late return/reminder letter was sent 6 weeks following the first mail-out. Data collection took place from March to August 2017. A total of 825 patients completed the questionnaire package (64.7% response rate).

Ethics

The study was approved by the Regional Committee for Medical Research Ethics for South East Norway (ID 2016/1745) and conducted in accordance with the Declaration of Helsinki. The study was also approved by the administrative heads of the Section for Climate Therapy at Oslo University Hospital and the Centre for Privacy and Information Security at Oslo University Hospital.

Measures

The HLQ is a widely used measure of HL that has been translated and used in many countries.³⁴ The HLQ includes 44

Table 1 Health Literacy Questionnaire (HLQ) scales with indication of number of items and range of response categories (Norwegian version)

HLQ scales	Number of items	Response categories
1 Feeling understood and supported by healthcare providers	4	1 = Strongly disagree; 2 = disagree;
2 Having sufficient information to manage my health	4	3 = agree; 4 = strongly agree
3 Actively managing my health	5	
4 Social support for health	5	
5 Appraisal of health information	5	
6 Ability to actively engage with healthcare providers	5	1 = Cannot do; 2 = very difficult; 3 = quite
7 Navigating the healthcare system	6	difficult; 4 = quite easy; 5 = very easy
8 Ability to find good health information	5	
9 Understanding health information well enough to know what to do	5	

items over nine independent scales, each representing a different element of the overall HL construct (Table 1). Each scale includes 4–6 items. The first five scales comprise items asking the respondents to indicate level of agreement [ranging from strongly disagree to strongly agree (range 1–4)]. The remaining scales (6–9) have domains of self-reported capability (range 1–5: ‘cannot do’ to ‘very easy’). The questionnaire has no total score, as that could potentially mask individual needs in specific HL domains.³⁴ The HLQ appears robust for its intended purpose of assessing HL in a range of settings and has shown excellent reliability.^{36–38}

Self-Administered Psoriasis Area and Severity Index

The Self-Administered Psoriasis Area and Severity Index (SAPASI) is a structured instrument that allows patients to assess accurately the severity of their psoriasis. The questionnaire uses the same criteria as the Psoriasis Area and Severity Index (PASI) but is presented using nonprofessional terminology.³⁹ SAPASI scores range from 0 to 72, where a higher score indicates more severe illness (Cronbach $\alpha = 0.74$).

Medical comorbidity is measured using an adapted and simplified version of the Self-Administered Comorbidity Questionnaire (SCQ-18),^{40,41} where higher scores indicate a more severe comorbidity profile.

The Psoriasis Knowledge Questionnaire (PKQ)⁴² assesses psoriasis knowledge based on 44 statements about psoriasis. The responses are reported as valid, uncertain or invalid, and the total calculated score range is 0–44, where higher scores indicate higher levels of knowledge (Cronbach $\alpha = 0.86$).

The Dermatology Life Quality Index (DLQI) measures quality of life (QoL). It contains 10 questions concerning symptoms and feelings, daily activities, personal relationships and treatment,⁴³ and is a frequently used QoL instrument in dermatology.⁴⁴ The scores for individual items (0–3) are added to yield a total score (0–30), where higher scores indicate greater impairment of a patient’s QoL (Cronbach $\alpha = 0.90$).

The General Self-Efficacy Scale assesses the beliefs that one can perform novel or difficult tasks in life or cope with hardship.⁴⁵ The scale has 10 items with a response range from 1 (not at all true) to 4 (exactly true). Correspondingly, sum

scores range from 10 to 40, where a higher score means higher self-efficacy (Cronbach $\alpha = 0.85$).

Statistical analysis

Descriptive statistics report characteristics of the study population. The expectation maximization algorithm was used to impute missing HLQ item scores.⁴⁶ For all HLQ scales, assumptions of normal distribution were met.

To investigate associations between variables, bivariate correlation analysis (Pearson’s *r*) and hierarchical linear multiple regression analysis (enter method) were used. In the hierarchical multiple regression analyses the nine HLQ scales were used as the independent variables, with four steps (presented in Table 2). In the results, the final step (step 4) is reported. Residuals had approximately normal distribution, and tests for linearity were not statistically significant, indicating that all associations were approximately linear. All statistical analyses were performed using SPSS version 25 (IBM, Armonk, NY, U.S.A.). *P*-values < 0.05 were considered statistically significant.

Results

Descriptive data

Patient demographics are shown in Table 3. The mean \pm SD age of the participants was 53.3 \pm 12.4 years (range 21–83);

Table 2 The independent variables added in the four steps of the regression model

Step 1	Age, sex, years living with psoriasis, educational attainment
Step 2	Plus additional diseases, number of climate heliotherapy treatments, disease severity (SAPASI) and quality of life (DLQI)
Step 3	Plus self-efficacy (GSES)
Step 4	Plus psoriasis knowledge (PKQ)

SAPASI, Self-Administered Psoriasis Area and Severity Index; DLQI, Dermatology Life Quality Index; GSES, General Self-Efficacy scale; PKQ, Psoriasis Knowledge Questionnaire.

Table 3 Demographic and clinical characteristics of Norwegian patients with psoriasis participating in climate helio therapy (CHT) from 2011 to 2016 (n = 825)

	N (%) / Mean \pm SD / Median (range)
Female sex	390 (47.3)
Age (years)	
Mean \pm SD	53.3 \pm 12.4
Range	18–83
Marital status	
Married/cohabiting	544 (65.9)
Unmarried/single	128 (15.5)
Divorced/separated/widowed	139 (16.9)
Others	12 (1.4)
Level of education (n = 822)	
Primary/secondary school \leq 10 years	94 (11.4)
Vocational/high school \leq 13 years	407 (49.5)
College/university < 3 years	182 (22.1)
College/university \geq 3 years	139 (16.9)
Median (range) duration of disease (years)	27 (0–77)
Mean \pm SD health condition (VAS scale 0–100)	60.11 \pm 19.6
Mean \pm SD self-assessed health status (1–5, i.e. poor–excellent)	3.33 \pm 0.92
Current smoker (n = 822)	193 (23.5)
Median (range) number of CHT treatments	2 (1–39)
Biological medicines (n = 811)	115 (14.2)
Joint pain	551 (66.8)
Joint pain and PsA affirmed by rheumatologist	254 (46.1)
Mean \pm SD BMI	28.64 \pm 5.30
Mean \pm SD number of comorbidities	4.4 \pm 2.50
Mean \pm SD SAPASI (0–72) ^a	7.51 \pm 4.93
Median (range) DLQI (0–30) ^b	9 (0–30)
Mean \pm SD PKQ (0–44) ^c	24.6 \pm 7.26
Mean \pm SD GSE (10–40) ^d	30.20 \pm 4.58

Data are n (%) unless otherwise indicated. VAS, visual analogue scale; PsA, psoriatic arthritis; BMI, body mass index; SAPASI, Self-Administered Psoriasis and Severity Index; DLQI, Dermatology Life Quality Index; PKQ, Psoriasis Knowledge Questionnaire; GSE, General Self-Efficacy scale. ^aHigher score indicates more serious disease; ^bhigher score indicates more impairment; ^chigher score indicates more knowledge; ^dhigher score indicates higher self-efficacy.

39% of participants were aged < 50 years. Women comprised 47.4% of the sample and 14.3% of the participants were currently receiving biological medicines. Previous histories of depression were noted in 35.5%, and 67.0% reported joint pain following psoriasis. The mean SAPASI and DLQI scores were 7.5 \pm 4.9 and 9.7 \pm 7.0, respectively. The participants reported a medium of 4.4 \pm 2.50 comorbidities (range 0–13).

The health literacy scales

The mean scores for each HLQ scale are shown in Table 4. The highest overall score was seen for scale 3, 'Actively

managing health' (mean score 2.78 \pm 0.51). The lowest score was for scale 5, 'Appraise health information' (mean score 2.54 \pm 0.54).

For scales 6–9, the highest and lowest scores were for scales 9, 'Ability to understand health information well enough to know what to do' (mean score 3.56 \pm 0.62) and 7, 'Ability to navigate the healthcare system' (mean score 3.10 \pm 0.71), respectively.

A total of 32 participants had missing data across one or more HLQ scales after the imputation procedure: no statistically significant differences were seen between those missing and not missing HLQ data, except that these 32 were significantly older (odds ratio –6.46, CI –11.35 to –1.57; *P* = 0.01).

Associations between health literacy domains and quality of life, demographic, clinical and self-management variables

The bivariate associations between demographic, clinical and self-management variables and QoL are presented in the first row of Table 5. The last step (i.e. step 4) in the analyses show the main results.

Regarding the demographic variables, age was not a significant predictor in any of the HL domains, and sex was only significant in the equation where scale 3, 'Actively managing my health', was the predicted variable. In this equation, men scored lower on actively managing their health (standardized β = 0.10). A higher educational level was a significant predictor for HL in scale 8, 'Ability to find good health information' (β = 0.13) and scale 9, 'Ability to understand health information well enough to know what to do' (β = 0.12).

Concerning the clinical variables, fewer years with psoriasis was significant with higher HL in scale 8, 'Ability to find good health information' (β = –0.08). Less comorbidity was significantly related to higher HL in scales 4, 'Having social support for health' (β = 0.12), and 7, 'Ability to navigate the healthcare system' (β = –0.10), whereas more comorbidity was related to higher HL in the first scale, 'Feeling understood and supported by healthcare providers' (β = 0.08). Higher disease severity measured by the SAPASI showed a significant relationship to higher HL in scale 9, 'Ability to understand health information' (β = 0.08), whereas numbers of climate helio treatments attended was not related to any of the HLQ scales.

Higher QoL measured by DLQI was a significant predictor of higher HL in seven of the nine HLQ scales (β = –0.13 to –0.23). Only scales 3, 'Actively managing health', and 5, 'Critical appraisal', were not related.

In terms of the self-management variables, higher self-efficacy showed a significant association with higher HL in all nine HLQ scales (β = 0.20–0.28). Regarding psoriasis knowledge, the PKQ was a significant predictor in seven of the scales; only scales 3, 'Actively managing health', and 4, 'Social support for health', were not related to higher psoriasis knowledge.

Table 4 Distribution of the Health Literacy Questionnaire scale scores (n = 825)

HLQ scales	Range	n	Mean ± SD
1 Feeling understood and supported by healthcare providers	1–4	804	2.72 ± 0.65
2 Having sufficient information to manage my health	1–4	804	2.64 ± 0.53
3 Actively managing my health	1–4	804	2.78 ± 0.51
4 Social support for health	1–4	805	2.55 ± 0.60
5 Appraisal of health information	1–4	804	2.54 ± 0.54
6 Ability to actively engage with healthcare providers	1–5	798	3.39 ± 0.73
7 Navigating the healthcare system	1–5	798	3.10 ± 0.71
8 Ability to find good health information	1–5	797	3.40 ± 0.65
9 Understanding health information enough to know what to do	1–5	797	3.56 ± 0.62

The fourth and final step in the linear regression model explained 9.6–22.1% of the variance in the HLQ scales (Table S1; see Supporting Information).

Discussion

This study provides the first step toward understanding the relationships connecting HL with sociodemographic and clinical variables, self-efficacy, knowledge and QoL in patients with psoriasis. The demographics showed that this large cohort of patients with psoriasis was middle-aged with a relatively long disease duration and a rather high comorbidity profile. They had a mean SAPASI score that indicates moderate illness severity, and their disease-related QoL indicates that their lives are considerably affected by psoriasis.⁴⁴

The results suggest that this population has weaknesses in HL. Their HLQ scores appear to be lower than those shown in other studies with different populations from other countries,^{47–49} and included other chronic diseases.^{23,50} It is concerning that a significant proportion of this population with psoriasis reports lower scores in so many domains of HL, especially given that this study was undertaken in Norway, which has a universal healthcare system with numerous strategies encouraging patient-centred care, patient participation and health communication. It seems particularly concerning in regard to the domains associated with understanding health information and navigating the healthcare system. This appears important as people with chronic conditions are increasingly expected to self-manage their own health.^{29,51}

A positive association between HL and disease severity (measured by SAPASI) was evident in scale 9, 'Ability to understand health information well enough to know what to do'. This seems somewhat counterintuitive as patients with severe psoriasis are likely to have had long-term engagement with specialist care and have received more education related to their treatment.⁵²

The results showed significant associations between more comorbidity and lower HL with regard to scales 4, 'Have social support for health', and 7, 'Ability to navigate the healthcare system'. Given that most information available to patients is based on single disease guidance, this finding is not surprising. People with multiple conditions may find that two

or more of their conditions have conflicting procedures and treatment recommendations,⁵³ and may require more support to self-manage. With only 24% of individuals in this study having fewer than three comorbid diseases, the problem of single disease guidelines presents a significant challenge to an individual's cognitive capacity, and their capability to be suitably health literate about their conditions. For instance, a large Danish study of people with other long-term conditions found that having more than one long-term condition was associated with more difficulties engaging with healthcare providers and understanding health information.²³ Several studies identified that patients with psoriasis do not receive follow-up tailored to living with a complex long-term disease,^{54,55} including a substantial cardiovascular burden.^{9,55} Previous psoriasis research highlights that patients experience difficulties in relation to healthcare providers, especially doctors.^{56–58} Concurrently, a rigid healthcare system with deficiency in communication across departments and specialties and different health service organization may also increase the burden for patients with multiple disorders and challenge HL.⁵⁹

QoL was strongly associated with seven of the HLQ scales throughout the regression analyses. Several studies have explored the relationship between HL and QoL, but the results seem inconclusive. For example HL has no impact on health-related QoL in patients with rheumatic diseases,⁶⁰ or among frequent users of healthcare services.⁶¹ In contrast, a large cohort study of general practices in the UK found that poor HL was associated with lower QoL scores (i.e. physical, psychological, social relationships and environment).⁶² In light of these conflicting results, it seems possible that the relationship between HL and QoL depends on certain aspects such as specific chronic diseases, cultural characteristics or other aspects of QoL. It seems plausible that interventions to strengthen HL in psoriasis may also have an additional positive impact on participants' QoL.

We found a significant association between HL and self-efficacy in all the HLQ scales. Knowing the value of self-efficacy to attain self-management this is an important finding that supports previous research.^{22,63–65} Patients with higher HL may feel more confident in their ability to pursue self-management behaviours and may be more likely to perform them.⁶⁶ Importantly one study that found self-efficacy to be the most

Table 5 Relationship between independent and dependent variables by hierarchal multiple regression analysis (n = 825) (r, standardized beta weights, adjusted R² and significance level)

Independent variables	(1) Feeling understood and supported by healthcare providers				(2) Having sufficient information to manage my health				(3) Actively managing health						
	r	Step 1 (β1)	Step 2 (β2)	Step 3 (β3)	Step 4 (β4)	r	Step 1 (β1)	Step 2 (β2)	Step 3 (β3)	Step 4 (β4)	r	Step 1 (β1)	Step 2 (β2)	Step 3 (β3)	Step 4 (β4)
Age (years)	.029	.01	-.02	-.02	.01	.04	.03	-.00	-.01	.04	.05	.05	.05	.05	.06
Sex (0 = men, 1 = women)	.016	-.00	.01	.02	.01	-.00	-.02	.01	.03	.01	.09**	.08*	.09*	.10*	.10**
Educational level 1–4 (higher score = higher level of education)	.068*	.09*	.08*	.03	-.00	.18***	.18***	.15***	.08*	.04	.15***	.15***	.14***	.08*	.07
Years with Psoriasis	.040	-.05	-.05	-.05	-.07	.056	-.01	-.01	-.01	-.04	.03	-.03	-.03	-.03	-.04
Quantity of comorbidities (higher score = more comorbidity)	-.002	.04	.08	.08	.08*	-.16***	-.12**	-.06	-.06	-.07	-.03	-.04	-.04	.01	.01
SAPASI (Higher score = more severe disease)	-.07*	.02	.02	-.01	.02	-.14***	-.06	-.07	-.07	-.06	-.05	-.07	-.07	-.04	-.03
Numbers of CHT (higher score = more climate treatments)	.11*	.12*	.12*	.11	.07	.10*	.12*	.10**	.10**	.06	.07	.08	.08	.06	.05
Quality of life (DLQI)	-.16***	-.18***	-.17***	-.17***	-.16***	-.21***	-.15**	-.14*	-.14*	-.13**	-.05	-.01	-.01	.00	.00
Self-efficacy (GSE) (higher score = better self-efficacy)	.22***	.21***	.21***	.21***	.20***	.34***	.29***	.29***	.29***	.28***	.26***	.26***	.25***	.25***	.24***
Psoriasis knowledge (PKQ) (Higher score = higher PSO knowledge)	.16***	.12**	.12**	.12**	.12**	.24***	.15***	.15***	.15***	.15***	.14***	.14***	.14***	.07	.05
Adjusted R ² change		.00	.03	.04***	.04***	.10	.03	.09	.16	.07***	.03	.03	.03	.08	.08

Independent variables	(4) Have social support for health				(5) Appraise health information				(6) Ability to actively engage with healthcare providers				
	Step 1 (β1)	Step 2 (β2)	Step 3 (β3)	Step 4 (β4)	Step 1 (β1)	Step 2 (β2)	Step 3 (β3)	Step 4 (β4)	r	Step 1 (β1)	Step 2 (β2)	Step 3 (β3)	Step 4 (β4)
Age (years)	.02	-.00	-.03	-.02	-.06*	-.07	-.07	-.01	.03	-.00	.01	.03	.03
Sex (0 = men, 1 = women)	.00	-.01	.05	.06	.03	.01	.03	-.01	-.06	-.04	-.02	-.07	-.03
Educational level	.07	.03	-.02	-.04	.14***	.14***	.08*	.02	.14***	.12**	.05	.15***	.02
1-4 (higher score = higher level of education)													
Years with Psoriasis	.04	.01	.03	-.002	-.03	-.04	-.04	-.08*	.049	.00	-.01	-.01	-.02
Quantity of comorbidities (higher score = more comorbidity)	-.19***	-.17***	-.15***	-.12**	-.02	.00	.06	.05	-.13***	-.08*	-.03	-.03	-.03
SAPASI (Higher score = more severe disease)	-.05	-.07	-.04	-.06	-.05	-.06	-.07	-.06	-.08*	-.03	-.02	-.02	.02
Numbers of CHT (higher score = more climate treatments)	.08	.09	.07	.06	.07	.10**	.08*	.02	.08	.08*	.065	.065	.04
Quality of life (DLQI)	-.19***	-.20***		-.19***	-.01		.04	.05	-.22***	-.21***	-.19***	-.19***	-.18***
Self-Efficacy (GSE) (higher score = better self-efficacy)	.29***		.26***	.25***	.29***	.02	.29***	.27***	.33***	.28***	.28***	.28***	.28***
Psoriasis knowledge (PKQ) (Higher score = higher PSO knowledge)	.12**		.05	.05	.27***			.23***	.17***				.17**
Adjusted R ² R ² change	.00	.07	.07***	.12	.02	.02	.10	.14	.02	.03***	.05***	.14	.15

Independent variables	(7) Ability to navigate the health care system				(8) Ability to find good health information				(9) Ability to understand health information well enough to know what to do				
	Step 1 (β1)	Step 2 (β2)	Step 3 (β3)	Step 4 (β4)	Step 1 (β1)	Step 2 (β2)	Step 3 (β3)	Step 4 (β4)	r	Step 1 (β1)	Step 2 (β2)	Step 3 (β3)	Step 4 (β4)
Age (years)	.04	-.01	-.02	.02	-.06	-.08	-.08	-.01	-.05*	-.05	-.05	-.05	.00
Sex (0 = men, 1 = women)	-.04	.00	.02	.00	-.002	-.00	.02	-.01	.02	-.01	-.05	-.05	.03
Educational level 1-4 (higher score = higher level of education)	.16***	.12**	.07	.04	.27***	.25***	.19***	.13**	.26***	.23***	.18***	.18***	.12**
Years with Psoriasis	.09**	.03	.03	.01	-.02	-.04	-.04	-.08*	-.02	-.05	-.06	-.06	-.09*
Quantity of comorbidities (higher score = more comorbidity)	-.19***	-.14***	-.010*	-.10***	-.14***	-.07	-.02	-.03	-.14***	-.09*	-.05	-.05	-.05
SAPASI (Higher score = more severe disease)	-.15***	-.01	-.021	-.02	-.12**	-.01	-.08	-.01	-.02	.08	.07	.07	.08*
Numbers of CHT (higher score = more climate treatments)	.11*	.12*	.10**	.07	.04	.08*	-.06	.00	.07	.10*	.08*	.08*	.02
Quality of life (DLQI)	-.29***	-.25***	-.24***	-.23***	-.21***	-.18***	-.17***	-.16***	-.14***	-.16**	-.14***	-.14***	-.13*
Self-Efficacy (GSE) (higher score = better self-efficacy)	.30***		.23***	.23***	.32***	.25***	.25***	.23***	.31***	.23***	.24***	.23***	.23***
Psoriasis knowledge (PKQ) (Higher score = higher PSO knowledge)	.20***			.13**	.33***			.26***	.29***				.22***
Adjusted R2	.03	.13.11***	.17.05***	.18.01*	.07	.11.05***	.17.06***	.22.05***	.06	.09.04***	.14.05***	.14.05***	.18.04***
R2 change													

Step1: Standardized beta weights using age, sex and educational attainment as independent variables. Step 2: Standardized beta weights using age, sex, educational attainment, years living with psoriasis, additional diseases, SAPASI, numbers of CHT and quality of life (DLQI) as independent variables. Step 3: Standardized beta weights using age, sex, educational attainment, years living with psoriasis, additional diseases, SAPASI, numbers of CHT, quality of life and self-efficacy (GSE) as independent variables. Step4: Standardized beta weights using age, sex, years living with psoriasis, educational attainment, additional diseases, SAPASI, numbers of CHT, quality of life, self-efficacy and Psoriasis Knowledge (PKQ) as independent variables. *Significant at the .05 level, **Significant at the .01 level, ***Significant at the .001 level.

important predictor of diabetes self-care also found that although HL was positively associated with self-efficacy, and self-efficacy was positively associated with self-care behaviours, the correlation between HL and self-care behaviours was statistically insignificant. These results indicate that HL may be a possible antecedent to self-efficacy and that the influence of HL on self-care behaviours may be mediated through self-efficacy.⁶⁷ This possible association needs more research.

Better psoriasis knowledge, measured by the PKQ, was strongly correlated with higher HL in seven of the HLQ scales and predicted a relevant R^2 change in all HL scales. Similar results have been found in research investigating a variety of disease conditions.^{67–70} Previous studies in psoriasis have demonstrated knowledge gaps concerning treatment and self-management,^{26,27,29} but the relationships to HL are not yet addressed. Findings from a study in patients with rheumatoid arthritis indicated that HL was independently associated with medication knowledge but not medication adherence.⁷¹ In a review exploring the impact of HL on self-management skills in 31 chronic disease studies, a consistent association was found between low HL and poorer disease-related knowledge in the case of respiratory diseases, diabetes and multiple disease categories.²² A previous study within the same psoriasis population has shown that being a woman, higher education, higher PASI score and previous CHT participation were significantly associated with greater psoriasis-related knowledge.⁴² It seems reasonable to expect that people with higher HL may be more aware of the importance of psoriasis knowledge to keep symptoms under control and also more prone to participate in patient education and self-management programmes. For patients with psoriasis with lower HL levels it may be difficult to adhere to complex treatment modalities and navigate lifestyle changes related to optimal self-management.¹⁴

Unlike the prevailing literature,^{47,72,73} we did not find that lower education was associated with lower health literacy, other than in scales 8, 'Ability to find good health information' and 9, 'Ability to understand health information well enough to know what to do.' As these scales indicate an ability to be an 'information explorer', using several informational sources and an ability to understand written information and adequately complete forms,⁴⁶ the relationship with educational level seems obvious. Van der Heide *et al.* found in their medication analysis that HL plays a larger role in those with less education than among those with more.⁷⁴

To our knowledge, the present study is the first to relate the HLQ to psoriasis in primary care. An additional strength is the large sample size and representativeness of the sample, as well as the use of the comprehensive and valid measure HLQ.⁷⁵

Next, a little over two-thirds of all eligible patients (65%) agreed to take part in this study. This response rate is high compared with the response rates for previous studies using similar methods within chronic care.^{62,76}

It is a limitation that HL was measured using a paper-based questionnaire, which, by design, excluded illiterate individuals. Additionally, we only provided the Norwegian language version. This may have reduced the response rate from people with

a non-Norwegian background. The study is also limited by the self-reported survey, without clinical verification of health conditions, such as the number of comorbidities or verified psoriatic arthritis, and these values could differ from information held in medical records. Another limitation is the somewhat limited sample. All participants had undergone CHT and even though their ages ranged from 18 to 83 years, they were, to a high degree, middle-aged. This may reduce the generalizability of the results.

Initiatives for strengthening the role of patients in health care may be improved by paying more attention to patients' HL. Health literacy enables people to build their knowledge, skills and potential to make positive behavioural changes and, for example, the teach-back method encourages patients to use their own words to describe what they have learned.⁷⁷

A systematic review found that very different intervention types and settings were associated with changes in health literacy (73% of all studies).⁷⁸ The review showed that both group and individual interventions of varying intensity in primary healthcare and community settings are useful in supporting sustained change in health literacy to foster change in behavioural risk factors. Unfortunately, HL interventions also tend to lack rigorous study design and therefore it may be premature to try to select the optimal interventions.⁷⁹ Improving health literacy training for healthcare professionals has received increasing emphasis in recent years.⁸⁰ However, the existing literature seems to be heavily skewed toward the field of medicine.⁸¹ In psoriasis, based on our findings, improving HL may be a useful strategy for reducing disparities in self-management skills, self-efficacy and QoL. However, it appears that more research is needed to uncover the complex relationships and factors that interfere in these mechanisms before we can tailor efficient interventions and improve health care.

Acknowledgments

We wish to thank the patients with psoriasis who participated in the study and the staff at the Section for Climate Therapy, Oslo University Hospital, for help with the questionnaire administration.

References

- Gibbs S. Skin disease and socioeconomic conditions in rural Africa: Tanzania. *Int J Dermatol* 1996; **35**:633–9.
- Danielsen K, Duvetorp A, Iversen L *et al.* Prevalence of psoriasis and psoriatic arthritis and patient perceptions of severity in Sweden, Norway and Denmark: results from the Nordic Patient Survey of Psoriasis and Psoriatic Arthritis. *Acta Derm Venereol* 2019; **99**:18–25.
- World Health Organization. *Global Report on Psoriasis Switzerland*. Geneva: Department for Management of Noncommunicable Diseases, Disability, Violence and Injury Prevention, 2016.
- Russo PA, Ilchef R, Cooper AJ. Psychiatric morbidity in psoriasis: a review. *Australas J Dermatol* 2004; **45**:155–61.
- Linder MD, Piaserico S, Augustin M *et al.* Psoriasis – the life course approach. *Acta Derm Venereol* 2016; **96**:102–8.

- 6 Kimball AB, Gieler U, Linder D et al. Psoriasis: is the impairment to a patient's life cumulative? *J Eur Acad Dermatol Venereol* 2010; **24**:989–1004.
- 7 Kimball AB, Guerin A, Tsaneva M et al. Economic burden of comorbidities in patients with psoriasis is substantial. *J Eur Acad Dermatol Venereol* 2011; **25**:157–63.
- 8 Ijosaa TM, Mork C, Stubhaug A et al. Skin pain and skin discomfort is associated with quality of life in patients with psoriasis. *J Eur Acad Dermatol Venereol* 2012; **26**:29–35.
- 9 Augustin M, Vietri J, Tian H, Gilleteau I. Incremental burden of cardiovascular comorbidity and psoriatic arthritis among adults with moderate-to-severe psoriasis in five European countries. *J Eur Acad Dermatol Venereol* 2017; **31**:1316–23.
- 10 Bundy C, Borthwick M, McAtteer H et al. Psoriasis: snapshots of the unspoken: using novel methods to explore patients' personal models of psoriasis and the impact on well-being. *Br J Dermatol* 2014; **171**:825–31.
- 11 Meyer N, Paul C, Feneron D et al. Psoriasis: an epidemiological evaluation of disease burden in 590 patients. *J Eur Acad Dermatol Venereol* 2010; **24**:1075–82.
- 12 Gottlieb AB, Chao C, Dann F. Psoriasis comorbidities. *J Dermatolog Treat* 2008; **19**:5–21.
- 13 Rodríguez-Zúñiga MJM, García-Perdomo HA. Systematic review and meta-analysis of the association between psoriasis and metabolic syndrome. *J Am Acad Dermatol* 2017; **77**:657–66.
- 14 Rasmussen GS, Maindal HT, Lomborg K. Self-management in daily life with psoriasis: an integrative review of patient needs for structured education. *Nurs Res Pract* 2012; **2012**:890860.
- 15 van Cranenburgh OD, de Korte J, Sprangers MA et al. Satisfaction with treatment among patients with psoriasis: a web-based survey study. *Br J Dermatol* 2013; **169**:398–405.
- 16 Tan J, Wolfe B. Improved decisional conflict and preparedness for decision making using a patient decision aid for treatment selection in psoriasis: a pilot study. *J Cutan Med Surg* 2014; **18**:114–18.
- 17 Buchbinder R, Batterham R, Ciciriello S et al. Health literacy: what is it and why is it important to measure? *J Rheumatol* 2011; **38**:1791–7.
- 18 Federman AD, Wolf MS, Sofianou A et al. Self-management behaviors in older adults with asthma: associations with health literacy. *J Am Geriatr Soc* 2014; **62**:872–9.
- 19 Souza JG, Apolinario D, Magaldi RM et al. Functional health literacy and glycaemic control in older adults with type 2 diabetes: a cross-sectional study. *BMJ Open* 2014; **4**:e004180.
- 20 Puente-Maestu L, Calle M, Rodríguez-Hermosa JL et al. Health literacy and health outcomes in chronic obstructive pulmonary disease. *Respir Med* 2016; **115**:78–82.
- 21 Gazmararian JA, Williams MV, Peel J Baker DW. Health literacy and knowledge of chronic disease. *Patient Educ Couns* 2003; **51**:267–75.
- 22 Mackey LM, Doody C, Werner EL, Fullen B. Self-management skills in chronic disease management: what role does health literacy have? *Med Decis Making* 2016; **36**:741–59.
- 23 Friis K, Lasgaard M, Osborne RH et al. Gaps in understanding health and engagement with healthcare providers across common long-term conditions: a population survey of health literacy in 29,473 Danish citizens. *BMJ Open* 2016; **6**:e009627.
- 24 Gutknecht M, Schaarschmidt ML, Herrlein O, Augustin M. A systematic review on methods used to evaluate patient preferences in psoriasis treatments. *J Eur Acad Dermatol Venereol* 2016; **30**:1454–64.
- 25 Conrad C, Gilliet M. Psoriasis: from pathogenesis to targeted therapies. *Clin Rev Allergy Immunol* 2018; **54**:102–113.
- 26 Renzi C, Di Pietro C, Gisondi P et al. Insufficient knowledge among psoriasis patients can represent a barrier to participation in decision-making. *Acta Derm Venereol* 2006; **86**:528–34.
- 27 Skiveren J, Philipsen P, Therning G. Patients with psoriasis have insufficient knowledge of their risk of atherothrombotic disease and metabolic syndrome. *Clin Exp Dermatol* 2015; **40**:600–4.
- 28 Lee MK, Kim HS, Cho EB et al. A study of awareness and screening behavior of cardiovascular risk factors in patients with psoriasis and dermatologists. *Ann Dermatol* 2015; **27**:59–65.
- 29 Rasmussen GS, Kragballe K, Maindal HT, Lomborg K. Experience of being young with psoriasis: self-management support needs. *Qual Health Res* 2017; **28**:73–86.
- 30 Nelson PA, Kane K, Chisholm A et al. 'I should have taken that further' – missed opportunities during cardiovascular risk assessment in patients with psoriasis in UK primary care settings: a mixed-methods study. *Health Expect* 2016; **19**:1121–37.
- 31 Blome C, von Usslar K, Augustin M. Feasibility of using qualitative interviews to explore patients' treatment goals: experience from dermatology. *Patient* 2016; **9**:261–9.
- 32 Zweegers J, van den Reek JMPA, van de Kerkhof PCM et al. Comparing treatment goals for psoriasis with treatment decisions in daily practice: results from a prospective cohort of patients with psoriasis treated with biologics: BioCAPTURE. *Br J Dermatol* 2014; **171**:1091–8.
- 33 Wahl AK, Langeland E, Larsen MH et al. Positive changes in self-management and disease severity following climate therapy in people with psoriasis. *Acta Derm Venereol* 2015; **95**:317–21.
- 34 Osborne RH, Batterham R, Elsworth G et al. The grounded psychometric development and initial validation of the Health Literacy Questionnaire (HLQ). *BMC Public Health* 2013; **13**:658.
- 35 Wahl AK, Mork C, Cooper BA, Padilla G. No long-term changes in psoriasis severity and quality of life following climate therapy. *J Am Acad Dermatol* 2005; **52**:699–701.
- 36 Maindal HT, Kayser L, Norgaard O et al. Cultural adaptation and validation of the Health Literacy Questionnaire (HLQ): robust nine-dimension Danish language confirmatory factor model. *Springerplus* 2016; **5**:1232.
- 37 Nolte S, Osborne RH, Dwinger S et al. German translation, cultural adaptation, and validation of the Health Literacy Questionnaire (HLQ). *PLOS ONE* 2017; **12**:e0172340.
- 38 Hawkins M, Elsworth GR, Osborne RH. Application of validity theory and methodology to patient-reported outcome measures (PROMs): building an argument for validity. *Qual Life Res* 2018; **27**:1695–710.
- 39 Sampogna F, Sera F, Mazzotti E et al. Performance of the self-administered psoriasis area and severity index in evaluating clinical and sociodemographic subgroups of patients with psoriasis. *Arch Dermatol* 2003; **139**:353–8.
- 40 Sangha O, Stucki G, Liang MH et al. The self-administered comorbidity questionnaire: a new method to assess comorbidity for clinical and health services research. *Arthritis Care Res* 2003; **49**:156–63.
- 41 Stolwijk C, van Tubergen A, Ramiro S et al. Aspects of validity of the self-administered comorbidity questionnaire in patients with ankylosing spondylitis. *Rheumatology* 2014; **53**:1054–64.
- 42 Wahl AK, Moum T, Robinson HS et al. Psoriasis patients' knowledge about the disease and treatments. *Dermatol Res Pract* 2013; **2013**:921737.
- 43 Finlay AY, Khan GK. Dermatology Life Quality Index (DLQI) – a simple practical measure for routine clinical use. *Clin Exp Derm* 1994; **19**:210–16.
- 44 Lewis V, Finlay AY. 10 years experience of the Dermatology Life Quality Index (DLQI). *J Invest Dermatol* 2004; **9**:169–80.
- 45 Luszczynska A, Scholz U, Schwarzer R. The General Self-Efficacy Scale: multicultural validation studies. *J Psych* 2005; **139**:439–57.
- 46 Dodson S, Beauchamp A, Batterham R, Osborne R. Ophelia Toolkit: A step-by-step guide for identifying and responding to health literacy needs within local communities. Available at: <https://www.>

- ophelia.net.au/bundles/opheliapublic/pdf/DEA1238-Ophelia-Toolkit-FA-Digital_reduced.pdf (last accessed 1 March 2018).
- 47 Jessup RL, Osborne RH, Beauchamp A *et al.* Health literacy of recently hospitalised patients: a cross-sectional survey using the Health Literacy Questionnaire (HLQ). *BMC Health Serv Res* 2017; **17**:52.
 - 48 Kolarcik P, Cepova E, Madarasova Geckova A *et al.* Structural properties and psychometric improvements of the Health Literacy Questionnaire in a Slovak population. *Int J Public Health* 2017; **62**:591–604.
 - 49 van der Gaag M, van der Heide I, Spreeuwenberg PMM *et al.* Health literacy and primary health care use of ethnic minorities in the Netherlands. *BMC Health Serv Res* 2017; **17**:350.
 - 50 Olesen K, Reynheim AL, Joensen L *et al.* Higher health literacy is associated with better glycemic control in adults with type 1 diabetes: a cohort study among 1399 Danes. *BMJ Open Diabetes Res Care* 2017; **5**:e000437.
 - 51 Jonkman NH, Schuurmans MJ, Jaarsma T *et al.* Self-management interventions: proposal and validation of a new operational definition. *J Clin Epidemiol* 2016; **80**:34–42.
 - 52 Al Sawah S, Foster SA, Goldblum OM *et al.* Healthcare costs in psoriasis and psoriasis sub-groups over time following psoriasis diagnosis. *J Med Econ* 2017; **20**:982–90.
 - 53 Allison W, Elizabeth M, Rowan W. Interventions to improve medication adherence in people with multiple chronic conditions: a systematic review. *J Adv Nurs* 2008; **63**:132–43.
 - 54 Chisholm A, Nelson PA, Pearce CJ *et al.* The role of personal models in clinical management: Exploring health care providers' beliefs about psoriasis. *Br J Health Psychol* 2016; **21**:114–34.
 - 55 Ersser SJ. Providing lifestyle behaviour change support for patients with psoriasis: an assessment of the existing training competencies across medical and nursing health professionals. *Br J Dermatol* 2014; **171**:450–1.
 - 56 Halioua B, Maury Le Breton A, de Fontaubert A *et al.* Treatment refusal among patients with psoriasis. *J Dermatolog Treat* 2015; **26**:396–400.
 - 57 Uhlenhake EE, Kurkowski D, Feldman SR. Conversations on psoriasis – what patients want and what physicians can provide: a qualitative look at patient and physician expectations. *J Dermatolog Treat* 2010; **21**:6–12.
 - 58 Magin PJ, Adams J, Heading GS, Pond CD. Patients with skin disease and their relationships with their doctors: a qualitative study of patients with acne, psoriasis and eczema. *Med J Aust* 2009; **190**:62–4.
 - 59 Mirzaei M, Aspin C, Essue B *et al.* A patient-centred approach to health service delivery: improving health outcomes for people with chronic illness. *BMC Health Serv Res* 2013; **13**:251.
 - 60 Zhang X-H, Li S-C, Fong K-Y, Thumboo J. The impact of health literacy on health-related quality of life (HRQoL) and utility assessment among patients with rheumatic diseases. *Value Health* 2009; **12**:106–S9.
 - 61 Couture ÉM, Chouinard M-C, Fortin M, Hudon C. The relationship between health literacy and quality of life among frequent users of health care services: a cross-sectional study. *Health Qual Life Outcomes* 2017; **15**:137.
 - 62 Panagioti M, Skevington SM, Hann M *et al.* Effect of health literacy on the quality of life of older patients with long-term conditions: a large cohort study in UK general practice. *Qual Life Res* 2018; **27**:1257–68.
 - 63 Bandura A. *Self-efficacy: The Exercise of Control*. New York: Freeman, 1997.
 - 64 Walton B, Vivienne WSF, Chieh-Yu L *et al.* Health literacy, self-efficacy, and self-care behaviors in patients with type 2 diabetes mellitus. *J Am Assoc Nurse Pract* 2013; **25**:495–502.
 - 65 Larsen MH, Krogstad AL, Aas E *et al.* A telephone-based motivational interviewing intervention has positive effects on psoriasis severity and self-management: a randomized controlled trial. *Br J Dermatol* 2014; **171**:1458–69.
 - 66 World Health Organization. *Health Education: Theoretical Concepts, Effective Strategies and Core Competencies: A Foundation Document to Guide Capacity Development of Health Educators*. Cairo: WHO Regional Office for the Eastern Mediterranean, 2012.
 - 67 Reisi M, Mostafavi F, Javadzade H *et al.* Impact of health literacy, self-efficacy, and outcome expectations on adherence to self-care behaviors in Iranians with type 2 diabetes. *Oman Med J* 2016; **31**:52–9.
 - 68 Williams MV, Baker DW, Parker RM, Nurss JR. Relationship of functional health literacy to patients' knowledge of their chronic disease: a study of patients with hypertension and diabetes. *Arch Intern Med* 1998; **158**:166–72.
 - 69 Bailey SC, Brega AG, Crutchfield TM *et al.* Update on health literacy and diabetes. *Diabetes Educ* 2014; **40**:581–604.
 - 70 Al Sayah F, Majumdar SR, Williams B *et al.* Health literacy and health outcomes in diabetes: a systematic review. *J Gen Intern Med* 2013; **28**:444–52.
 - 71 Quinlan P, Price KO, Magid SK *et al.* The relationship among health literacy, health knowledge, and adherence to treatment in patients with rheumatoid arthritis. *HSS J* 2013; **9**:42–9.
 - 72 Juul L, Rowlands G, Maingal HT. Relationships between health literacy, motivation and diet and physical activity in people with type 2 diabetes participating in peer-led support groups. *Prim Care Diabetes* 2018; **12**:331–7.
 - 73 van der Heide I, Poureslami I, Mitic W *et al.* Health literacy in chronic disease management: a matter of interaction. *J Clin Epidemiol* 2018; **102**:134–8.
 - 74 van der Heide I, Wang J, Droomers M *et al.* The relationship between health, education, and health literacy: results from the Dutch Adult Literacy and Life Skills Survey. *J Health Commun* 2013; **18**(Suppl. 1):172–84.
 - 75 Elsworth GE, Beauchamp A, Osborne RH. Measuring health literacy in community agencies: a Bayesian study of the factor structure and measurement invariance of the Health Literacy Questionnaire (HLQ). *BMC Health Serv Res* 2016; **16**:508.
 - 76 Guthrie EA, Dickens C, Blakemore A *et al.* Depression predicts future emergency hospital admissions in primary care patients with chronic physical illness. *J Psychosom Res* 2016; **82**:54–61.
 - 77 Schillinger D, Piette J, Grumbach K *et al.* Closing the loop: physician communication with diabetic patients who have low health literacy. *Arch Intern Med* 2003; **163**:83–90.
 - 78 Taggart J, Williams A, Dennis S *et al.* A systematic review of interventions in primary care to improve health literacy for chronic disease behavioral risk factors. *BMC Fam Pract* 2012; **13**:49.
 - 79 Brainard J, Loke Y, Salter C *et al.* Healthy ageing in Europe: prioritizing interventions to improve health literacy. *BMC Res Notes* 2016; **9**:270.
 - 80 Coleman CA, Hudson S, Maine LL. Health literacy practices and educational competencies for health professionals: a consensus study. *J Health Commun* 2013; **18**(Suppl. 1):82–102.
 - 81 Coleman C. Teaching health care professionals about health literacy: a review of the literature. *Nurs Outlook* 2011; **59**:70–8.

Supporting Information

Additional Supporting Information may be found in the online version of this article at the publisher's website:

Powerpoint S1. Journal Club Slide Set.